

## ANURA

**BUFFER** - How much riparian buffer or wooded area does the species need around its breeding site?

Species	Common Name	Buffer
<b>Bufonidae</b>	<b>True Frogs</b>	
<i>Bufo a. americanus</i>	Eastern American Toad	22.8 - 475.5 m forested (Oldham 1966)
<i>Bufo fowleri</i>	Fowler's Toad	
<i>Bufo quercicus</i>	Oak Toad	Unk
<i>Bufo terrestris</i>	Southern Toad	Unk
<b>Hylidae</b>	<b>Treefrogs</b>	
<i>Acris c. crepitans</i>	Eastern Cricket Frog	Unk, but remain in the vicinity of the [breeding] waters throughout the year (Hulse et al. 2001)
<i>Acris c. blanchardi</i>	Blanchard's Cricket Frog	
<i>Acris g. gryllus</i>	Coastal Plain Cricket Frog	
<i>Hyla andersonii</i>	Pine Barrens Treefrog	
<i>Hyla chrysoscelis</i>	Cope's Gray Treefrog	Unk (Little 1983)
<i>Hyla cinerea</i>	Green Treefrog	
<i>Hyla femoralis</i>	Pine Woods Treefrog	Unk
<i>Hyla gratiosa</i>	Barking Treefrog	Unk
<i>Hyla squirella</i>	Squirrel Treefrog	
<i>Hyla versicolor</i>	Gray Treefrog	Unk
<i>Pseudacris brachyphona</i>	Mountain Chorus Frog	Unk
<i>Pseudacris brimleyi</i>	Brimley's Chorus Frog	Unk
<i>Pseudacris c. crucifer</i>	Northern Spring Peeper	Unk
<i>Pseudacris feriarum</i>	Upland Chorus Frog	Unk
<i>Pseudacris feriarum kalmi</i>	New Jersey Chorus Frog	Unk
<i>Pseudacris ocularis</i>	Little Grass Frog	Unk
<i>Pseudacris triseriata</i>	Western Chorus Frog	Unk
<b>Pelobatidae</b>	<b>Spadefoot Toads</b>	
<i>Scaphiopus holbrookii</i>	Eastern Spadefoot	Unk
<b>Ranidae</b>	<b>True Frogs</b>	
<i>Rana catesbeiana</i>	American Bullfrog	Unk
<i>Rana clamitans melanota</i>	Northern Green Frog	Unk
<i>Rana palustris</i>	Pickerel Frog	Unk
<i>Rana pipiens</i>	Northern Leopard Frog	Unk
<i>Rana septentrionalis</i>	Mink Frog	Unk
<i>Rana sphenoccephala</i>	Southern Leopard Frog	Unk
<i>Rana sylvatica</i>	Wood Frog	Unk
<i>Rana virgatipes</i>	Carpenter Frog	Unk
<b>Microhylidae</b>	<b>Narrow-mouthed Frogs</b>	
<i>Gastrophryne carolinensis</i>	Eastern Narrow-mouthed Toad	Unk

**Buffer Codes:** buffer distance (e.g., 50 m) or Unk = unknown